



## Note

## Rafting in Zoantharia: a hitchhiker's guide to dispersal?

Maria E.A. Santos<sup>a,\*</sup>, James Davis Reimer<sup>a,b</sup><sup>a</sup> Molecular Invertebrate Systematics and Ecology Laboratory, Graduate School of Engineering and Science, University of the Ryukyus, Senbaru 1, Nishihara, Okinawa 903-0213, Japan<sup>b</sup> Tropical Biosphere Research Center, University of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan

## ARTICLE INFO

## Keywords:

Artificial substrates  
Biofouling  
Debris  
Human-made structures

## ABSTRACT

The increasing availability of human-made structure in the oceans coupled with climate changes may lead to the range expansion of species able to disperse by rafting. In this study, we report on zoantharian species of genera *Isaurus*, *Palythoa*, *Umimayanthus* and *Zoanthus* covering artificial substrates in locations in both the Atlantic and Pacific oceans. Moreover, we reviewed observations of additional zoantharians as macrofouling organisms, and discuss the possible role of rafting in the dispersal of this cosmopolitan group. Traits reported to some zoantharian species, such as hermaphroditism and resistance to desiccation, support their dispersal potential by drifting attached to floating objects. Further reports of zoantharian species covering floating artificial structures and natural debris are needed to increase our knowledge of dispersal mechanisms in the oceans. Additionally, this information is essential to monitor and manage possible exotic species invasions, especially for zoantharian species that are common in the aquarium trade.

## 1. Introduction

Rafting has been reported in many marine taxa that are able to disperse by attaching to floating drifting objects (Thiel and Gutow, 2005), for example, reef fishes with seaweed flotsam (Luiz et al., 2015), and sea anemones and hydrocorals attached to volcanic pumice (Jokiel, 1990; Bryan et al., 2012). Artificial floating structures, such as plastic and glass litter, are nowadays much more available than natural debris, in addition to having slower decomposition times (Derraik, 2002; Kiessling et al., 2015; Carlton et al., 2017). As a consequence, rafting on human-made structures has led to the introduction of non-native species and become a major environmental problem (Barnes, 2002; Kiessling et al., 2015; Rech et al., 2016, 2018).

Zoantharians (Cnidaria: Anthozoa: Zoantharia) are benthic animals found in all oceans (Haddon and Shackleton, 1891; Burnett et al., 1997; Ryland et al., 2000). Species of genera *Palythoa* and *Zoanthus* are a common component of shallow-water subtropical and tropical reefs and can cover large portions of the substrate (Karlson, 1983; Oigman-pszczol et al., 2004; Irei et al., 2011). This widespread distribution is likely associated with a high dispersal capability, due to a long pelagic larval duration (Ryland et al., 2000; Polak et al., 2011) and several asexual reproduction strategies, including polyp ball production (Acosta et al., 2005) and relatively rapid growth of fragments (Karlson, 1988). In addition, there is the possible role of rafting, although this potential has not been examined or discussed for zoantharians.

In this study, we report on zoantharian colonies covering portions of artificial substrates (bottle, mooring line, shipwreck). Furthermore, we link these reports to previous biofouling records of species of Zoantharia, and discuss the potential for dispersal of zoantharians via rafting.

## 2. Methods

## 2.1. Specimens on artificial substrates

Zoantharian polyps were observed growing on a glass fragment of a bottle in October 2014, during an expedition to Rocas Atoll (3° 86'S, 33° 80' W), 270 km off northeast Brazil, in the Atlantic Ocean. The atoll is approximately 3 km in diameter (Gherardi and Bosence, 2001) and the glass piece was collected from a sandy beach during low tide, where it had washed up.

In addition, zoantharian colonies were noticed covering large portions of the sunken Japanese tanker *Iro* (7°17'57"N, 134°24'56"E), and on a mooring line in Turtle Cove (7°05'13"N, 134°15'33"E), during field work in the Republic of Palau between 2016 and 2018. Zoantharian specimens were identified via examination of external morphology (Pax, 1910; Reimer, 2010; Reimer et al., 2012a; Montenegro et al., 2015). We also compiled previous published biofouling records of zoantharian species.

\* Corresponding author.

E-mail address: [santos.mariaea@gmail.com](mailto:santos.mariaea@gmail.com) (M.E.A. Santos).




Fig. 1. Glass fragment covered with *Palythoa grandiflora* and *Zoanthus sociatus* polyps collected from Rocas Atoll, northeast Brazil. The fragment was approximately 10 cm in length and had polyps on both external (A) and internal (B) sides.

### 3. Results

Colonies of the species *Palythoa grandiflora* (family Sphenopidae) and *Zoanthus sociatus* (family Zoanthidae) were found on a glass fragment from Rocas Atoll (Fig. 1A-B). The fragment was from a bottle and was approximately 10 cm in length, partially covered on both sides by approximately 35 zoantharian polyps. It is unknown if the species settled after or before the glass was broken. Moreover, several colonies of *Zoanthus sansibaricus* were observed on the external wall of the sunken tanker *Iro* in Palau, as well as numerous polyps of *Umimayanthus chanpuru* (family Parazoanthidae). In addition, a colony of *Isaurus tuberculatus* (family Zoanthidae) and *Palythoa cf. mutuki* were also noted in a mooring line in Palau. The line was made of nylon and was in midwater, attached to a fixed float. The zoantharian polyps were observed growing on surfaces already covered by other encrusting organisms such as encrusting coralline algae, both on the bottle fragment from Rocas and on the artificial substrates in Palau. Moreover, species of *Umimayanthus* have an obligatory epizoic relationship with Porifera hosts, and *U. chanpuru* colonies were associated with encrusting sponges.

Additionally, species of *Palythoa*, *Zoanthus* and *Microzoanthus* (family Microzoanthidae) have been noted as biofouling in previous literature (Table 1). Genera *Isaurus*, *Palythoa* and *Zoanthus* belong to suborder Brachycnemina, while *Microzoanthus* and *Umimayanthus* belong to suborder Macrocnemina.

### 4. Discussion

We observed zoantharian colonies attached to a glass surface, a plastic mooring line, and an artificial iron reef. The polyps were growing upon other encrusting organisms, which may have facilitated their initial settlement. Cases of zoantharians covering artificial structures are not rare (Table 1), and previous reports from the Atlantic Ocean include colonies of *Palythoa caribaeorum* on a wreck (Vieira et al., 2012), on concrete blocks (Brandini and Silva, 2011), and on reefs (Silva et al., 2016), as well as *P. mammosa* on a limestone boulder (Cummings, 1994). In the Indo-Pacific, species of *Palythoa* and *Zoanthus* have been noted as macrofouling organisms on jetty pilings in Singapore (Ong and Tan, 2012), and *Zoanthus* has been reported on a fiberglass experimental panel in the Persian Gulf (Naser, 2017). Furthermore, polyps of the zoantharian *Microzoanthus kagareou* were observed attached to sea water outflow pipes of an aquaculture center (Reimer and Fujii, 2013). There is no information available to

Table 1

Records of zoantharian species growing on artificial substrates.

Artificial substrate type	Species	Location(s)	Reference(s)
Concrete	<i>Palythoa caribaeorum</i>	Brazil, Mexico	Brandini and Silva, 2011; Silva et al., 2016
Fiberglass	<i>Zoanthus</i> sp.	Bahrain	Naser, 2017
Glass	<i>Palythoa grandiflora</i> <i>Zoanthus sociatus</i>	Brazil	This study
Jetty pilings	<i>Palythoa</i> sp. <i>Zoanthus</i> sp.	Singapore Singapore	Ong and Tan, 2012 Ong and Tan, 2012
Limestone boulder	<i>Palythoa mammosa</i>	USA	Cummings, 1994
Nylon mooring line	<i>Isaurus tuberculatus</i> <i>Palythoa cf. mutuki</i>	Palau Palau	This study This study
Shipwreck	<i>Palythoa caribaeorum</i> <i>Umimayanthus chanpuru</i> <i>Zoanthus sansibaricus</i>	Brazil Palau Palau	Vieira et al. 2012 This study This study
Outflow pipes	<i>Microzoanthus kagareou</i>	Palau	Reimer and Fujii, 2013

determine in these above cases if polyps were growing upon other organisms or not, and such data should be included in future observations. In summary, the current and previous reports demonstrate the ability of zoantharians to settle on substrates of several types and sizes, likely including floating objects. It is possible that zoantharians are also able to grow on natural debris, similar to as seen in other benthic cnidarians such as sea anemones (Bryan et al., 2012), hydrocorals (Jokiel, 1989, 1990), and scleractinian corals (Bryan et al., 2004, 2012), that have been reported on volcanic sea-rafting pumice. Moreover, the scleractinian coral *Astrangia poculata* has been reported to cross long distances on metal flotsam (Hoeksema et al., 2012, 2015). Likewise, genetic data indicate that rafting may be the main dispersal mode of the ascidian *Symplesma rubra*, which can be found on both natural and anthropogenic floating objects (Dias et al., 2006). Rafting may also drive gene flow among long distant areas for some marine invertebrates (Nikula et al., 2012), which could be a key dispersal strategy in species that do not have a long pelagic larval stage. For instance, the wide distribution and the occurrence of *Millepora alcicornis* in remote areas is probably due to successful raft dispersal, as although this hydrocoral may have a short pelagic phase, it shows generalist

settlement behavior (De Souza et al., 2017; Hoeksema et al., 2017).

The ability to grow on different artificial substrates includes representative species of both suborders of Zoantharia (Brachyneremina and Macrocnemina), and reports are from several locations in different oceans. Species settlements on glass, metal and nylon surfaces (Table 1) indicate that these zoantharians could potentially grow and raft on objects such as detached metal flotsam, mooring line or fishing glass buoys. Traits reported for some *Palythoa* and *Zoanthus* species such as relatively high resistance against desiccation (Sebens, 1982; Rabelo et al., 2015), hermaphroditism (Fadlallah et al., 1984), and high frequency of fertile polyps throughout the year (Boscolo and Silveira, 2005) also lend support to high dispersal potential by rafting. Similar to some other marine groups, the capability of zoantharians to attach to different structures may enable them to colonize and disperse across oceans attached to floating substrates, and may be especially important for species with shorter pelagic larvae durations. However, zoantharian species occurring in environments such as caves (Irei et al., 2015) or sandy areas (Reimer et al., 2012b) are unlikely to be good candidates for rafting. It is estimated that some rafting species are not yet known (Rech et al., 2018), and Zoantharia is a group that is often overlooked in marine surveys (Burnett et al., 1995; Santos et al., 2016). Therefore, reports of zoantharians on floating objects of both natural and artificial origins could increase our knowledge of the dispersal mechanisms and routes of marine species. Moreover, rafting can turn into a significant environmental issue with the invasion of alien species associated to artificial floating objects such as plastic and glass bottles, in addition to secondary dispersal of already-introduced species (Rech et al., 2016; Gracia et al., 2018). The growing numbers of human-made structures in the oceans, in addition to an increasing occurrence of large storms (e.g. typhoons) and global ocean warming, may lead to distributional range expansions of rafting species (Hoeksema et al., 2012; Carlton et al., 2017). Thus, macrofouling reports of Zoantharia are needed to help in predicting and monitoring potential exotic species invasions (Creed et al., 2017). Such reports may be particularly important due to the popularity of some *Palythoa* and *Zoanthus* species in the global aquarium trade (Deeds et al., 2011), and an associated higher risk of invasive introductions (Whitfield et al., 2002). Further investigations of zoantharian species' rafting abilities in addition to research on their reproduction and growth rates will help to better understand the processes behind Zoantharia zoogeography, as well as of possible species' range expansions.

## 5. Conclusions

Species from at least five genera of Zoantharia are able to grow on artificial substrates, such as glass and nylon, and this capability likely includes floating objects. Additionally, some *Palythoa* and *Zoanthus* species have traits that support their potential to raft across long distances. Similar to other benthic cnidarians, zoantharians may also grow on natural floating objects, and confirmation of this hypothesis will help to better understand zoantharian species' distributions. Further observations are essential to increase our knowledge of Zoantharia dispersal mechanisms, as well as to monitor possible alien species.

## Acknowledgements

We thank two anonymous reviewers for comments that improved the quality of this manuscript. Research in Rocas Atoll was supported by Rocas Atoll Reserve/ICMBio and SISBIOTA-Mar network (CNPq 563276/2010-0 and FAPESC 6308/2011-8). We are grateful to “Zelinha” de Brito Silva for her contribution to field work in Rocas. Work in Palau was supported by the SATREPS P-CORIE Project “Sustainable management of coral reef and island ecosystem: responding to the threat of climate change”, funded by the Japan Science and Technology Agency and the Japan International Cooperation Agency in cooperation with Palau International Coral Reef Center

(PICRC) and Palau Community College. Hiroki Kise (U. Ryukyus, PICRC), Pat Colin and Lori Colin (Coral Reef Research Foundation) are also thanked for information and field assistance in Palau. MEAS was additionally supported by a Japanese Government (MEXT) scholarship, and JDR by a JSPS ‘Kiban B’ grant entitled ‘Global evolution of Brachyneremina and their *Symbiodinium*’.

## Author contribution

MEAS performed field observations, analyzed the data, prepared figures, conceived and wrote the paper. JDR performed field observations, analyzed the data, reviewed drafts of the paper, and provided funding for the majority of the field work. Both authors read and approved the final manuscript.

## Conflict of interest

None.

## References

- Acosta, A., Sammarco, P.W., Duarte, L.F., 2005. New fission processes in the zoanthid *Palythoa caribaeorum*: description and quantitative aspects. *Bull. Mar. Sci.* 76, 1–26.
- Barnes, D.K.A., 2002. Invasions by marine life on plastic debris. *Nature* 416, 808–809. <http://dx.doi.org/10.1038/416808a>.
- Boscolo, H.K., Silveira, F.L., 2005. Reproductive biology of *Palythoa caribaeorum* and *Protospalythoa variabilis* (Cnidaria, Anthozoa, Zoanthidea) from the southeastern coast of Brazil. *Reprod. Biol.* 65, 29–41.
- Brandini, F., Silva, A., 2011. Epilithic community development on artificial reefs deployed along a cross-shelf environmental gradient off Paraná state, southern Brazil. *Braz. J. Oceanogr.* 59, 43–53. <http://dx.doi.org/10.1590/S1679-87592011000500007>.
- Bryan, S.E., Cook, A.G., Evans, J.P., et al., 2004. Pumice rafting and faunal dispersion during 2001–2002 in the Southwest Pacific: record of a dacitic submarine explosive eruption from Tonga. *Earth Planet. Sci. Lett.* 227, 135–154. <http://dx.doi.org/10.1016/j.epsl.2004.08.009>.
- Bryan, S.E., Cook, A.G., Evans, J.P., et al., 2012. Rapid, long-distance dispersal by pumice rafting. *PLoS One* 7 (7), 1–13.
- Burnett, J., Benzie, J., Beardmore, J., Ryland, J., 1995. Patterns of genetic subdivision in populations of a clonal cnidarian, *Zoanthus coppingeri*, from the Great Barrier Reef. *Mar. Biol.* 122, 665–673.
- Burnett, W.J., Benzie, J.A.H., Beardmore, J.A., Ryland, J.S., 1997. Zoanthids (Anthozoa, Hexacorallia) from the Great Barrier Reef and Torres Strait, Australia: systematics, evolution and a key to species. *Coral Reefs* 16, 55–68.
- Carlton, J.T., Chapman, J.W., Geller, J.B., et al., 2017. Tsunami-driven rafting: transoceanic species dispersal and implications for marine biogeography. *Science* 356, 1402–1406.
- Creed, J.C., Fenner, D., Sammarco, P., et al., 2017. The invasion of the azooxanthellate coral *Tubastraea* (Scleractinia: Dendrophylliidae) throughout the world: history, pathways and vectors. *Biol. Invasions* 19, 283–305. <http://dx.doi.org/10.1007/s10530-016-1279-y>.
- Cummings, S.L., 1994. Colonization of a nearshore artificial reef at Boca Raton (Palm Beach County), Florida. *Bull. Mar. Sci.* 55, 1193–1215.
- De Souza, J.N., Nunes, F.L.D., Zilberberg, C., Sanchez, J.A., Migotto, A.E., Hoeksema, B.W., Serrano, X.M., Baker, A.C., Lindner, A., 2017. Contrasting patterns of connectivity among endemic and widespread fire coral species (*Millepora* spp.) in the tropical Southwestern Atlantic. *Coral Reefs* 36, 701–716. <http://dx.doi.org/10.1007/s00338-017-1562-0>.
- Deeds, J.R., Handy, S.M., White, K.D., Reimer, J.D., 2011. Palytoxin found in *Palythoa* sp. zoanthids (Anthozoa, Hexacorallia) sold in the home aquarium trade. *PLoS One* 6, 2–10. <http://dx.doi.org/10.1371/journal.pone.0018235>.
- Derraik, J.G.B., 2002. The pollution of the marine environment by plastic debris: a review. *Mar. Pollut. Bull.* 44, 842–852. [http://dx.doi.org/10.1016/S0025-326X\(02\)00220-5](http://dx.doi.org/10.1016/S0025-326X(02)00220-5).
- Dias, G.M., Lembo Duarte, L.F., Solferini, V.N., 2006. Low genetic differentiation between isolated populations of the colonial ascidian *Sympygmia rubra* Monniot, C. 1972. *Mar. Biol.* 148, 807–815. <http://dx.doi.org/10.1007/s00227-005-0111-5>.
- Fadlallah, Y.H., Karlson, R.H., Sebens, K.P., 1984. A comparative study of sexual reproduction in three species of Panamanian zoanthids (Coelenterata: Anthozoa). *Bull. Mar. Sci.* 35, 80–89.
- Gherardi, D.F.M., Bosence, D.W.J., 2001. Composition and community structure of the coralline algal reefs from Atol das Rocas, South Atlantic, Brazil. *Coral Reefs* 19, 205–219. <http://dx.doi.org/10.1007/s003380000100>.
- Gracia, A.C., Rangel-Buitrago, N., Flórez, P., 2018. Beach litter and woody-debris colonizers on the Atlantic department Caribbean coastline, Colombia. *Mar. Pollut. Bull.* 128, 185–196.
- Haddon, A.C., Shackleton, A.M., 1891. The Zoantheae. In: *A Revision of the British Actiniae*. Part II. Reports on the Zoological Collections Made in the Torres Straits by A. C. Haddon, 1888–1889. *Sci. Trans. R. Dublin Soc. Ser. 2* 4(12). pp. 609–672.

- Hoeksema, B.W., Roos, P.J., Cadee, G.C., 2012. Trans-Atlantic rafting by the brooding reef coral *Favia fragum* on man-made flotsam. *Mar. Ecol. Prog. Ser.* 445, 209–218. <http://dx.doi.org/10.3354/meps09460>.
- Hoeksema, B.W., Nunes, F.L.D., Lindner, A., De Souza, J.N., 2017. *Millepora alcicornis* (Hydrozoa: Capitata) at Ascension Island: confirmed identity based on morphological and molecular analyses. *J. Mar. Biol. Assoc. UK* 97, 709–712.
- Hoeksema, B.W., Ross, P.J., Cadee, G.C., 2015. Corrigendum to: trans-Atlantic rafting by the brooding reef coral *Favia fragum* on man-made flotsam. *Mar. Ecol. Prog. Ser.* 541, 279.
- Irei, Y., Nozawa, Y., Reimer, J.D., 2011. Distribution patterns of five zoanthid species at Okinawa Island, Japan. *Zool. Stud.* 50, 426–433.
- Irei, Y., Sinniger, F., Reimer, J.D., 2015. Descriptions of two azooxanthellate *Palythoa* species (subclass Hexacorallia, order Zoantharia) from the Ryukyu Archipelago, southern Japan. *Zookeys* 26, 1–26.
- Jokiel, P.L., 1989. Rafting of reef corals and other organisms at Kwajalein Atoll. *Mar. Biol.* 101, 483–493.
- Jokiel, P.L., 1990. Long-distance dispersal by rafting: reemergence of an old hypothesis. *Endeavour* 14, 66–73.
- Karlson, R.H., 1983. Disturbance and monopolization of a spatial resource by *Zoanthus sociatus* (Coelenterata, Anthozoa). *Bull. Mar. Sci.* 13, 118–131.
- Karlson, R.H., 1988. Size-dependent growth in two zoanthid species: a contrast in clonal strategies. *Ecology* 69, 1219–1232.
- Kiessling, T., Gutow, L., Thiel, M., 2015. Marine litter as habitat and dispersal vector. In: Bergmann, M., Gutow, L., Klages, M. (Eds.), *Marine Anthropogenic Litter*. Springer, Cham.
- Luiz, O.J., Allen, A.P., Robertson, D.R., et al., 2015. Seafarers or castaways: ecological traits associated with rafting dispersal in tropical reef fishes. *J. Biogeogr.* 41, 2323–2333. <http://dx.doi.org/10.1111/jbi.12574>.
- Montenegro, J., Sinniger, F., Reimer, J.D., 2015. Unexpected diversity and new species in the sponge-Parazoanthidae association in southern Japan. *Mol. Phylogenet. Evol.* 89, 73–90.
- Naser, H.A., 2017. Variability of marine macrofouling assemblages in a marina and a mariculture centre in Bahrainian gulf. *Reg. Stud. Mar. Sci.* 16, 162–170.
- Nikula, R., Spencer, H.G., Waters, J.M., 2012. Passive rafting is a powerful driver of transoceanic gene flow. *Biol. Lett.* 9, 1–4. <http://dx.doi.org/10.1098/rsbl.2012.0821>.
- Oigman-pszczol, S.S., Figueiredo, M.A.D.O., Creed, J.C., 2004. Distribution of benthic communities on the tropical rocky subtidal of Armação dos Búzios, Southeastern Brazil. *Mar. Ecol.* 25, 173–190.
- Ong, J.L.J., Tan, K.S., 2012. Observations on the subtidal fouling community on jetty pilings in the Southern Islands of Singapore. *Contrib. Mar. Sci.* 121–126.
- Pax, F., 1910. Studien an westindischen Actinien. *Zool. Jahrb. Abt. Anat. Ontog. Tiere* (Suppl. 11), 157–330.
- Polak, O., Loya, Y., Brickner, I., 2011. The widely-distributed Indo-Pacific zoanthid *Palythoa tuberculosa*: a sexually conservative strategist. *Bull. Mar. Sci.* 87, 605–621.
- Rabelo, E.F., Rocha, L.L., Colares, G.B., et al., 2015. *Symbiodinium* diversity associated with zoanthids (Cnidaria: Hexacorallia) in northeastern Brazil. *Symbiosis* 64, 105–113. <http://dx.doi.org/10.1007/s13199-014-0308-9>.
- Rech, S., Borrell Pichs, Y.J., García-Vazquez, E., 2016. Marine litter as a vector for non-native species: what we need to know. *Mar. Pollut. Bull.* 113, 40–43. <http://dx.doi.org/10.1016/j.marpolbul.2016.08.032>.
- Rech, S., Borrell Pichs, Y.J., Garcia-Vasquez, E., 2018. Anthropogenic marine litter composition in coastal areas may be a prediction of potentially invasive rafting fauna. *PLoS One* 13 (1), e0191859.
- Reimer, J.D., 2010. Key to field identification of shallow water brachycnemid zoanthids (order Zoantharia: suborder Brachycnemina) present in Okinawa. *Galaxea* 12, 23–29. <http://dx.doi.org/10.3755/galaxea.12.23>.
- Reimer, J.D., Fujii, T., 2013. First record of Microzoanthidae (Anthozoa: Hexacorallia: Zoantharia) in Palau and as a biofouling organism. *Mar. Biodivers. Rec.* 6, e103. <http://dx.doi.org/10.1017/S1755267213000808>.
- Reimer, J.D., Foord, C., Irei, Y., 2012a. Species diversity of shallow water zoanthids (Cnidaria: Anthozoa: Hexacorallia) in Florida. *J. Mar. Biotechnol.* <http://dx.doi.org/10.1155/2012/856079>.
- Reimer, J.D., Lin, M., Fujii, T., Lane, D.J.W., Hoeksema, B.W., 2012b. The phylogenetic position of the solitary zoanthid genus *Sphenopus* (Cnidaria: Hexacorallia). *Contrib. Zool.* 81, 43–54.
- Ryland, J.S., De Putron, S., Scheltema, R.S., et al., 2000. *Semper's* (zoanthid) larvae: pelagic life, parentage and other problems. *Hydrobiologia* 440, 191–198.
- Santos, M.E.A., Kitahara, M.V., Lindner, A., Reimer, J.D., 2016. Overview of the order Zoantharia (Cnidaria: Anthozoa) in Brazil. *Mar. Biodivers.* 46, 547–559. <http://dx.doi.org/10.1007/s12526-015-0396-7>.
- Sebens, K.P., 1982. Intertidal distribution of zoanthids on the Caribbean coast of Panamá: effects of predation and desiccation. *Bull. Mar. Sci.* 32, 316–335.
- Silva, R., Mendoza, E., Mariño-Tapia, I., et al., 2016. An artificial reef improves coastal protection and provides a base for coral recovery. *J. Coast. Res.* 75, 467–471. <http://dx.doi.org/10.2112/SI75-094.1>.
- Thiel, M., Gutow, L., 2005. The ecology of rafting in the marine environment. I. The floating substrata. *Oceanogr. Mar. Biol. Annu. Rev.* 42, 181–263. <http://dx.doi.org/10.1201/9780203507810.ch6>.
- Whitfield, P.E., Gardner, T., Vives, S.P., Gilligan, M.R., Courtenay Jr., W.R., Ray, G.C., Hare, J.A., 2002. Biological invasion of the Indo-Pacific lionfish *Pterois volitans* along the Atlantic coast of North America. *Mar. Ecol. Prog. Ser.* 235, 289–297.
- Vieira, L., Farrapeira, C., Amaral, F.D., Lira, S., 2012. Bryozoan biodiversity in Saint Peter and Saint Paul Archipelago, Brazil. *Cah. Biol. Mar.* 53, 159–167.